



(Invited Talk)

## Orthogonal Polynomials Related to a Complex Moment Functional: Theory and Applications

Gradimir V. Milovanović<sup>(1),(2)</sup>

<sup>(1)</sup>*Serbian Academy of Sciences and Arts, 11000 Belgrade, Serbia*  
*e-mail: gvm@mi.sanu.ac.rs*

<sup>(2)</sup>*University of Niš, Faculty of Sciences and Mathematics, P.O. Box 224, 18000 Niš, Serbia*  
*e-mail: gradimir.milovanovic@pmf.ni.ac.rs*

### Abstract

Motivated by certain applications in physics, in [4] the numerical calculation of the integrals  $\int_{-1}^1 f(x)W(x; \alpha, \beta)dx$ , with complex parameters in the Jacobi weight function, under the assumption  $\operatorname{Re}(\alpha) > -1$  and  $\operatorname{Re}(\beta) > -1$ , was considered. However, these applications in theoretical physics (calculation of the so-called transition amplitude [5, 6]) require further weakening of the conditions to the level of  $\operatorname{Re}(\beta) = -1$ . Let us mention that for complex parameters there are other types of orthogonality for Jacobi polynomials, such as the one in [3].

In order to obtain an adequate method for the aforementioned application, in this lecture we introduce orthogonal polynomials with respect to the linear complex-valued moment functional  $L[z^k] = \mu_k$ ,  $k = 0, 1, \dots$ , defined in a suitable way (cf. [1], [2], [7]). Beside the existence and uniqueness we prove several properties, including zero distribution. In the second part we construct the corresponding quadrature formulas of Gaussian and anti-Gaussian type.

**Keywords:** Complex moment functional, orthogonal polynomial, moments, weight function, recurrence relation, zero distribution, Gaussian quadrature formula, nodes, Christoffel numbers.

### References:

- [1] T. S. Chihara, *An Introduction to Orthogonal Polynomials*. Gordon & Breach, New York, 1978.
- [2] W. Gautschi, *Orthogonal Polynomials: Computation and Approximation*. Oxford University Press, Oxford, 2004.
- [3] A. B. J. Kuijlaars, A. Martínez-Finkelshtein and R. Orive, Orthogonality of Jacobi polynomials with general parameters. *Electron. Trans. Numer. Anal.* 19 (2005), 1–17.
- [4] G. V. Milovanović and A. S. Cvetković, Numerical integration with complex Jacobi weight function. In: S. Margenov, L. G. Vulkov and J. Waśniewski (eds.) *Numerical Analysis and Its Applications (4th International Conference, NAA 2008 Lozenetz, Bulgaria, June 16–20, 2008, Revised Selected Papers)*, pp. 20–31, Springer-Verlag, Berlin–Heidelberg, 2009.
- [5] I. Mančev, Continuum distorted wave – Born initial state (CDW – BIS) model for single charge exchange. *J. Comput. Meth. Sci. & Engineer.* 5 (2005), 73–89.
- [6] I. Mančev, N. Milojević and Dž. Belkić, Boundary-corrected four-body continuum-intermediate-state method: Single-electron capture from heliumlike atomic systems by fast nuclei, *Phys. Rev. A* 91 (2015), 062705, 20 pp.
- [7] G. Mastroianni and Milovanović, *Interpolation Processes – Basic Theory and Applications*, Springer Monographs in Mathematics, Springer-Verlag, Berlin–Heidelberg, 2008.